

Decoherence from Isocurvature Perturbations in Inflation

Tomislav Prokopec and Gerasimos I. Rigopoulos*
*Institute for Theoretical Physics, Utrecht University,
 Postbus 80.195, 3508 TD Utrecht, The Netherlands*

We discuss the possible role of isocurvature perturbations for the quantum decoherence of the curvature perturbation during inflation. We point out that if the inflaton trajectory in field space is curved, the adiabatic mode is generically coupled to the isocurvature mode and thus tracing out the latter can cause the curvature perturbation to decohere. We explicitly investigate this suggestion in a model of inflation with two decoupled massive fields and find that decoherence is effective for a wide range of mass ratios. In particular, we calculate the entanglement entropy for this model and show that it grows after horizon exit, providing a quantitative measure for decoherence.

I. INTRODUCTION

Inflation has become the leading paradigm of early universe cosmology not least because of its ability to imprint scale invariant inhomogeneities on superhorizon scales via a causal mechanism [1]. These inhomogeneities are thought to provide the seeds which later become the temperature anisotropies in the Cosmic Microwave Background and the Large Scale Structure in the Universe. In inflation, the origin of these initial perturbations lies in quantum fluctuations of matter fields, amplified during a period of quasi-exponential expansion, which source adiabatic and isocurvature cosmological perturbations. The adiabatic perturbation is a variation in total energy density or equivalently in the gravitational potential, while isocurvature perturbations are variations in the entropy density of various matter components. All inflationary models share this basic mechanism, with different models characterized by small deviations from exact scale invariance. Along with other calculable properties, these deviations are particularly appealing as they allow for inflationary models to be tested against increasingly precise observations [2].

Inflationary calculations are based on the quantum mechanics of scalar fields in expanding spacetimes [3], where the relevant observable is the amplitude of the field's Fourier modes. Although treated as a quantum mechanical variable, this amplitude is interpreted as a stochastic random variable described by a gaussian distribution, with the variance given by the power-spectrum. This interpretation, used in CMB analyses and simulations of Large Scale Structure, proves to be very accurate for calculational purposes; quantum correlation functions, exactly calculated, are practically the same as the corresponding correlation functions obtained by using the relevant classical probability distribution [4].

The consistency of this stochastic interpretation requires a density matrix which is diagonal in the amplitude basis. However, the density matrix of inflationary perturbations is *not* automatically diagonal in this basis (see eg. [5]). One cannot assign a specific amplitude to the Fourier modes which must thus be considered to exist in coherent superpositions of different amplitudes. Prior to a stochastic interpretation, an as-of-yet undisclosed decoherence mechanism is required to diagonalize the density matrix.

The phenomenon of decoherence is ubiquitous in quantum physics and originates from couplings of the system of interest with degrees of freedom belonging to some unobservable environment [6, 7]. Various arguments and calculations suggesting that a form of such environmental decoherence can indeed occur for inflationary perturbations have been put forward in [8, 9, 10, 11, 12, 13, 14, 15, 16]. In [9, 10, 11, 12, 13, 14, 16], the environment is taken to consist of the short wavelength modes which are coupled to the long wavelength modes via non-linear couplings. In this paper, we point out that the decoherence of cosmological perturbations can occur in a precisely defined dynamical setting if the role of the environment is played by unobservable isocurvature perturbations generated during multi-field inflation. The coupling in this case are bilinear and no non-linearities are invoked.

II. QUANTUM MECHANICS OF INFLATIONARY PERTURBATIONS

To study decoherence in inflation it is convenient to use the Schrödinger picture which we briefly review in this section. The action for a free massive scalar field is

$$S = -\frac{1}{2} \int d^4x \sqrt{-g} (\partial_\mu \phi \partial^\mu \phi + m^2 \phi^2) \quad (1)$$

* Current address: Helsinki Institute of Physics, P.O.Box 64, FIN-00014, University of Helsinki, Finland

where we take the spacetime metric to be

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -dt^2 + a(t)^2 \delta_{ij} dx^i dx^j. \quad (2)$$

Then, the canonical momentum conjugate to the field ϕ ,

$$\pi = \frac{\partial \mathcal{L}}{\partial(\partial_t \phi)} = a^3 \partial_t \phi, \quad (3)$$

allows us to write the Hamiltonian $H \equiv \int d^3x (\pi \partial_t \phi - \mathcal{L})$

$$H = \int d^3x \frac{1}{2} \left(\frac{\pi^2}{a^3} + a(\partial_i \phi)^2 + a^3 m^2 \phi^2 \right). \quad (4)$$

Quantization dictates the replacement $\pi(\mathbf{x}) \rightarrow -i\hbar \frac{\delta}{\delta \phi(\mathbf{x})}$ which realizes the commutation relation $[\phi(\mathbf{x}), \pi(\mathbf{y})] = i\hbar \delta(\mathbf{x} - \mathbf{y})$. The wave functional $\Psi[\phi]$ obeys the Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} \Psi = \hat{H} \Psi. \quad (5)$$

Since we will be considering perturbations in their ground states, we take the wavefunctional to have a gaussian form

$$\Psi[\phi] = \mathcal{N} \exp \left[-\frac{1}{2} \int d^3x d^3y \phi(\mathbf{x}) A(\mathbf{x}, \mathbf{y}, t) \phi(\mathbf{y}) \right], \quad (6)$$

with $\mathcal{N}$ the normalization factor. Due to the homogeneity of the vacuum state, the correlator in (6) satisfies, $A(\mathbf{x}, \mathbf{y}, t) = A(\mathbf{y}, \mathbf{x}, t)$. For this state the Schrödinger equation gives

$$i\hbar \partial_t \ln \mathcal{N} = \frac{\hbar^2}{2a^3} \int d^3x A(\mathbf{x}, \mathbf{x}, t) \quad (7)$$

$$i\hbar \partial_t A(\mathbf{y}, \mathbf{z}, t) = \frac{\hbar^2}{a^3} \int d^3x A(\mathbf{x}, \mathbf{y}, t) A(\mathbf{x}, \mathbf{z}, t) + a(\partial_{\mathbf{y}}^2 - a^2 m^2) \delta^{(3)}(\mathbf{y} - \mathbf{z}). \quad (8)$$

Since the theory is noninteracting, it is convenient to solve (8) in momentum space, where it becomes local. Indeed, upon writing

$$\phi(\mathbf{x}) = \int \frac{d^3k}{(2\pi)^3} \phi_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{x}}, \quad (9)$$

$$A(\mathbf{x}, \mathbf{y}, t) = \int \frac{d^3k}{(2\pi)^3} A(\mathbf{k}, t) e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})}, \quad (10)$$

with $\phi_{-\mathbf{k}} = \phi_{\mathbf{k}}^*$ and $A(-\mathbf{k}, t) = A(\mathbf{k}, t)$, we get

$$i\hbar \partial_t A(\mathbf{k}, t) = \frac{\hbar^2}{a^3} A^2(\mathbf{k}, t) - a(k^2 + a^2 m^2). \quad (11)$$

The *Heisenberg picture* mode functions $\psi_{\mathbf{k}}(\eta)$ of a scalar field with the action (1) satisfy

$$\partial_\eta^2 (a\psi_{\mathbf{k}}(\eta)) + \left(k^2 + \frac{\frac{1}{4} - \nu^2}{\eta^2} \right) (a\psi_{\mathbf{k}}(\eta)) = 0, \quad (12)$$

where η is the conformal time, defined via $a d\eta = dt$, and $\nu^2 = \frac{9}{4} - \frac{m^2}{H^2}$. The Bunch-Davies vacuum solution of (12) is of the form

$$a\psi_{\mathbf{k}}(\eta) = \sqrt{-\frac{\pi\eta}{4}} e^{i\frac{\pi}{2}(\nu + \frac{1}{2})} H_\nu^{(1)}(-k\eta), \quad (13)$$

and the correlator $A(\mathbf{k})$ (11) is related to the mode functions (13) via [3]

$$A(\mathbf{k}, \eta) = \frac{1}{2\hbar |\psi_{\mathbf{k}}(\eta)|^2} (1 - i a^2 \partial_\eta |\psi_{\mathbf{k}}(\eta)|^2). \quad (14)$$

At early times, in the regime $k^2\eta^2 \gg 1$ when the mode is “deep inside the horizon” and spacetime curvature is not important, the solution (13) or equivalently (6) and (14) reduces to that of Minkowski vacuum. At late times, when $k^2\eta^2 \ll 1$, the mode has “exited the horizon” and amplification (“particle production”) occurs. The wave function for each mode can be written as¹

$$\Psi(\phi_{\mathbf{k}}, \eta) \propto \exp\left(-\frac{1}{2}\phi_{\mathbf{k}}A(\mathbf{k}, \eta)\phi_{\mathbf{k}}^*\right), \quad (15)$$

where the normalization factor can be easily obtained from (7) and by requiring $\int d\phi d\phi^* |\Psi|^2 = 1$. From now on we shall be ignoring such factors. Note that in the Schrödinger picture the time evolution of the wave function is determined by the correlator $A(\mathbf{k}, \eta)$; $\phi_{\mathbf{k}}$ is *time independent* and should not be confused with the Heisenberg picture mode $\psi_{\mathbf{k}}(\eta)$ (13).

The quantum mechanics of cosmological perturbations has been studied in [3, 17]. The system evolves into a kind of state known as a squeezed state, which exhibits a high degree of WKB classicality

$$\hat{p}\Psi \simeq \partial_q S(q)\Psi, \quad (16)$$

where $\hat{q}$ and $\hat{p}$ are the configuration variable and its conjugate momentum, while S is the exponent of the wave function, practically the classical action. Thus, the amplitude of each mode and its conjugate momentum are related to a very high degree of accuracy via the corresponding classical relation. In cosmology, this state of affairs is interpreted as equivalent to a statistical mixture of mode amplitude eigenstates, where the probability that any of these states has been realized in our universe is given by $|\Psi(\phi_{\mathbf{k}}, \eta)|^2$. This would correspond to a density matrix which is diagonal in the field amplitude basis

$$\hat{\rho}_{\mathbf{k}} = \sum_{\phi_{\mathbf{k}}} |\Psi(\phi_{\mathbf{k}}, \eta)|^2 |\phi_{\mathbf{k}}\rangle\langle\phi_{\mathbf{k}}|. \quad (17)$$

Such a stochastic mixture is a very good approximation when calculating correlators such as $\langle\phi_{\mathbf{k}}^2\rangle$ [4], which are then related to the stochastic properties of cosmological perturbations. However, the wave function (15) does not directly lead to the interpretation suggested by (17). Explicitly calculating the density matrix from (15), one finds

$$\langle\phi_{\mathbf{k}}|\hat{\rho}_{\mathbf{k}}|\bar{\phi}_{\mathbf{k}}\rangle \equiv \rho(\phi_{\mathbf{k}}, \bar{\phi}_{\mathbf{k}}) = \Psi(\phi_{\mathbf{k}}, \eta)\Psi^*(\bar{\phi}_{\mathbf{k}}, \eta) \propto \exp\left(-u_{\mathbf{k}}\Re[A]u_{\mathbf{k}}^* - \frac{1}{4}\Delta_{\mathbf{k}}\Re[A]\Delta_{\mathbf{k}}^* - i\Im[A]\Re[u_{\mathbf{k}}\Delta_{\mathbf{k}}^*]\right),$$

where we have defined $u = (\phi + \bar{\phi})/2$ and $\Delta = \phi - \bar{\phi}$. The above expression does not vanish for $\Delta \neq 0$, making the off-diagonal terms of the density matrix non-zero. Thus, strictly speaking, the description (17) is not valid and one cannot say that any particular mode amplitude has been realized with a certain probability. The modes of cosmological perturbations exist in coherent superpositions with different mode amplitudes. In order to diagonalize the density matrix, some process of decoherence must take place. As we show below, the existence of fields other than the inflaton during inflation can decohere the density matrix of cosmological perturbations even if no direct coupling between the different fields is assumed. The relevant coupling is ensured by the existence of scalar metric perturbations.

III. ISOCURVATURE PERTURBATIONS AND DECOHERENCE

A. Adiabatic and isocurvature perturbations

The existence of isocurvature modes is ubiquitous when more than one scalar fields are relevant during inflation. We will therefore consider a two-field model of inflation with a potential $V(\varphi, \chi)$. For cosmological perturbations for the metric and the scalar fields

$$\begin{aligned} \phi(\mathbf{x}, t) &= \phi(t) + \delta\phi(\mathbf{x}, t), & \chi(\mathbf{x}, t) &= \chi(t) + \delta\chi(\mathbf{x}, t), \\ ds^2 &= -(1 + 2\Phi(\mathbf{x}, t))dt^2 + a^2(t)(1 - 2\Phi(\mathbf{x}, t))d\mathbf{x}^2, \end{aligned} \quad (18)$$

¹ The reality of the field $\phi(\mathbf{x})$ implies that the modes $\phi_{-\mathbf{k}}$ and $\phi_{\mathbf{k}}$ are related as $\phi_{-\mathbf{k}} = \phi_{\mathbf{k}}^*$, which necessitates that wave functions be represented in terms of two-mode states [17].

the Lagrangian describing the dynamics of gauge invariant perturbations can be written as [18]

$$L = \int d^3x \frac{1}{2} \partial_\eta \mathbf{q} \partial_\eta \mathbf{q} - \frac{1}{2} \mathbf{q} (-\nabla^2 + (aH)^2 \Omega) \mathbf{q}, \quad (19)$$

where

$$\mathbf{q} = a(\delta\varphi + \frac{\dot{\varphi}}{H}\Phi)\hat{\mathbf{e}}_\varphi + a(\delta\chi + \frac{\dot{\chi}}{H}\Phi)\hat{\mathbf{e}}_\chi \equiv q_\varphi \hat{\mathbf{e}}_\varphi + q_\chi \hat{\mathbf{e}}_\chi, \quad (20)$$

with $\hat{\mathbf{e}}_\varphi$ and $\hat{\mathbf{e}}_\chi$ the unit basis vectors in the $\varphi - \chi$ field coordinates. The matrix Ω is given by

$$\Omega = \frac{\partial_a \partial_b V}{H^2} \mathbf{e}_a \otimes \mathbf{e}_b - (2 - \epsilon) \mathbf{1} - 2\epsilon \left((3 + \epsilon) \mathbf{e}_1 \otimes \mathbf{e}_1 + 2\eta^\parallel \mathbf{e}_1 \otimes \mathbf{e}_1 + \eta^\perp (\mathbf{e}_1 \otimes \mathbf{e}_2 + \mathbf{e}_2 \otimes \mathbf{e}_1) \right), \quad (21)$$

where $\epsilon = -\dot{H}/H^2$ and

$$\eta^\parallel \equiv -3 - \frac{\partial \mathbf{V} \cdot \mathbf{e}_1}{H|\dot{\phi}|}, \quad \eta^\perp \equiv -\frac{\partial \mathbf{V} \cdot \mathbf{e}_2}{H|\dot{\phi}|}, \quad (22)$$

are two slow roll parameters related to the acceleration parallel and perpendicular to the field velocity [18]. The subscripts $\{a, b\}$ take the values $\{\varphi, \chi\}$ and we have used the unit vectors $\hat{\mathbf{e}}_1$ and $\hat{\mathbf{e}}_2$ to be defined shortly. Note that even if scalar fields are decoupled and the potential term is diagonal, the q_a fields are still coupled due to the inclusion of the metric perturbation Φ . Thus, perturbations are always coupled gravitationally. As we discuss below this coupling can suffice to decohere the perturbations if one degree of freedom is traced out.

In order to proceed, it is convenient to define two new basis vectors, tangential and normal to the field trajectory [18, 19]:

$$\mathbf{e}_1 = \cos\theta \mathbf{e}_\varphi + \sin\theta \mathbf{e}_\chi \equiv \frac{\dot{\varphi}}{\sqrt{\dot{\varphi}^2 + \dot{\chi}^2}} \mathbf{e}_\varphi + \frac{\dot{\chi}}{\sqrt{\dot{\varphi}^2 + \dot{\chi}^2}} \mathbf{e}_\chi \quad (23)$$

$$\mathbf{e}_2 = -\sin\theta \mathbf{e}_\varphi + \cos\theta \mathbf{e}_\chi, \quad (24)$$

and express the Lagrangian in terms of components in this basis - we have

$$L = \int d^3x \frac{1}{2} (\partial_\eta \mathbf{q} + \mathbf{Zq})^\top (\partial_\eta \mathbf{q} + \mathbf{Zq}) - \frac{1}{2} \mathbf{q}^\top (-\nabla^2 + (aH)^2 \Omega) \mathbf{q}, \quad (25)$$

where $(m, n = 1, 2)$

$$\mathbf{q} = \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}, \quad \mathbf{q}_m = \mathbf{e}_m \cdot \mathbf{q}, \quad (26)$$

$$\Omega_{mn} = \mathbf{e}_m \Omega \mathbf{e}_n \quad (27)$$

is the matrix Ω expressed in the $\{\hat{\mathbf{e}}_1, \hat{\mathbf{e}}_2\}$ basis and

$$\mathbf{Z} = \partial_\eta \theta \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \quad (28)$$

Then, using the conjugate momentum $\pi(\mathbf{x}) = \frac{\partial \mathcal{L}}{\partial(\partial_\eta q(\mathbf{x}))}$ the Hamiltonian can be written as

$$H = \int d^3x \frac{1}{2} \pi^\top \pi - \frac{1}{2} \pi^\top \mathbf{Zq} - \frac{1}{2} \mathbf{q}^\top \mathbf{Z}^\top \pi + \frac{1}{2} \mathbf{q}^\top (-\nabla^2 + (aH)^2 \Omega) \mathbf{q}. \quad (29)$$

The quantum Hamiltonian is obtained via the replacement $\pi(\mathbf{x}) \rightarrow -i\hbar \frac{\delta}{\delta q(\mathbf{x})}$ and the wavefunction satisfies the Schrödinger equation

$$i\hbar \Psi = \hat{H} \Psi, \quad (30)$$

which, along with the vacuum ansatz

$$\Psi = N \exp \left(-\frac{1}{2} \int d^3x d^3y \mathbf{q}^\top(\mathbf{x}) \mathbf{B}(\mathbf{x} - \mathbf{y}) \mathbf{q}(\mathbf{y}) \right), \quad (31)$$

leads to

$$i\hbar\partial_\eta \mathbf{B}(\mathbf{x} - \mathbf{y}) = \hbar^2 \left(\int d^3z \mathbf{B}(\mathbf{x} - \mathbf{z}) \mathbf{B}(\mathbf{z} - \mathbf{y}) \right) + i\hbar [\mathbf{B}(\mathbf{x} - \mathbf{y}), \mathbf{Z}] + (\nabla_{\mathbf{x}}^2 - (aH)^2 \Omega) \delta^{(3)}(\mathbf{x} - \mathbf{y}), \quad (32)$$

$$i\hbar\partial_\eta \ln N = \frac{\hbar}{2} \int d^3x \mathbf{B}(0). \quad (33)$$

The solution to these equations provides the full quantum description of cosmological perturbations in the two-field system.

B. The reduced density matrix

The process of decoherence relies on the existence of degrees of freedom which couple to - get entangled with - the system of interest but which are not directly accessible. It is assumed that they cannot be measured along with the system and thus provide an uncontrollable environment. Tracing them out leads to the diagonalization of the density matrix for the system of interest and thus explains the appearance of a classical world where different alternatives - like Schrödinger's cat being dead and alive - do not interfere and are perceived as distinct [6, 7]. In Cosmology, the role of the cat is played by the long wavelength fluctuations which lead to structure formation.

In what follows we suggest that decoherence can occur for cosmological perturbations if the isocurvature component q_2 is considered to be the unobservable environment and is traced out. In cosmological considerations the two degrees of freedom in $\mathbf{q}$ describing perturbations are not directly observable. A more tangible remnant of inflation is the curvature perturbation which persists in later epochs of cosmic evolution and is the reason for structure formation. More precisely, it is given by a scalar quantity $\mathcal{R}$ which parameterizes the spatial curvature of comoving time slices (see e.g. [20]). During inflation it is related to q_1 by

$$\hat{\mathcal{R}} = \frac{H}{\sqrt{\dot{\phi}^2 + \dot{\chi}^2}} \frac{\hat{q}_1}{a}, \quad (34)$$

and for single field inflation it is constant on long wavelengths. The presence of an isocurvature perturbation makes $\mathcal{R}$ evolve in time [18, 19]

$$\frac{d}{dN} \hat{\mathcal{R}} = 2 \frac{\partial V \cdot \mathbf{e}_2}{\dot{\phi}^2 + \dot{\chi}^2} \frac{\hat{q}_2}{a}. \quad (35)$$

The hat acts as a reminder that we are dealing with quantum operators.

Any process which couples to $\hat{\mathcal{R}}$ during inflation is effectively measuring $\hat{q}_1$, the adiabatic component of the perturbations. A measurement of $\hat{q}_2$ via gravitational interactions alone would entail a coupling to $\frac{d}{dN} \hat{\mathcal{R}}$ during inflation as can be seen from (35). Thus, the gravitational perturbation $\hat{\mathcal{R}}$, which does not have its own independent dynamics, acts as an apparatus which “measures” - gets entangled with - $\hat{q}_1$. Let us now assume that after inflation both ϕ and χ decay into relativistic particles, generating no entropy perturbation and making $\hat{\mathcal{R}}$ the only remnant from the inflationary epoch.² In this case, information about $\frac{d}{dN} \hat{\mathcal{R}}$ during inflation is lost and all post-inflationary perturbations can be obtained from $\hat{\mathcal{R}}$ at the end of inflation, which is conserved on super-Hubble scales after $\hat{q}_2$ decays. Once perturbations reenter the horizon at late times, $\frac{d}{dN} \hat{\mathcal{R}}$ becomes different from zero again. This late time behaviour reflects late time dynamics of matter fields on small scales and has nothing to do with the inflationary $\frac{d}{dN} \hat{\mathcal{R}}$. Consequently, as information about $\hat{q}_2$ is lost, we can consider it to be an inaccessible environment which cannot be directly observed and can therefore be traced out of the density matrix.

Under the above assumption, the density matrix relevant for the adiabatic perturbation alone is the one obtained by tracing out q_2

$$\tilde{\rho}(q_1, \bar{q}_1) = \int dq_2 dq_2^* \Psi(q_1, q_2) \Psi^*(\bar{q}_1, q_2), \quad (36)$$

where the wavefunction Ψ is given by (31). Expressing the result in terms of the variables $u = (q_1 + \bar{q}_1)/2$ and $\Delta = q_1 - \bar{q}_1$, we find

$$\tilde{\rho}(u, \Delta) = \exp \left[-\frac{1}{2}(u, \Delta) \mathbf{C} \begin{pmatrix} u^* \\ \Delta^* \end{pmatrix} \right], \quad (37)$$

² This reasoning is not valid if a gravitational potential is generated after inflation via a curvaton type mechanism [21, 22].

with

$$\mathbf{C} \equiv \begin{pmatrix} 2 \left(\Re[\mathbf{B}_{11}] - \frac{\Re[\mathbf{B}_{12}]^2}{\Re[\mathbf{B}_{22}]} \right) & i \left(\Im[\mathbf{B}_{11}] - \frac{\Re[\mathbf{B}_{12}]\Im[\mathbf{B}_{22}]}{2\Re[\mathbf{B}_{22}]} \right) \\ i \left(\Im[\mathbf{B}_{11}] - \frac{\Re[\mathbf{B}_{12}]\Im[\mathbf{B}_{22}]}{2\Re[\mathbf{B}_{22}]} \right) & \frac{1}{2} \left(\Re[\mathbf{B}_{11}] + \frac{\Im[\mathbf{B}_{12}]^2}{\Re[\mathbf{B}_{22}]} \right) \end{pmatrix}. \quad (38)$$

As we remarked above, decoherence would imply that the Δ components are suppressed when $|\Delta| \neq 0$. This will happen if

$$C_{11}/C_{22} \ll 1, \quad (39)$$

since in this case the relative importance of the Δ over the u terms will diminish, making quantum correlations unimportant. Thus, we expect that decoherence will become effective if

$$\Im[\mathbf{B}_{12}] > 2\sqrt{\Re[\mathbf{B}_{11}]\Re[\mathbf{B}_{22}]}. \quad (40)$$

In inflationary models with two scalar fields, Ω_{12} in (32) is in general non zero, $\Omega_{12} \neq 0$ and thus $\mathbf{B}_{12} \neq 0$ (see also (43) and (46)). In particular, for inflationary perturbations and on superhorizon scales, the imaginary part $\Im[B_{12}]$ strongly dominates over the real parts $\Re[B_{11}]$ and $\Re[B_{22}]$ and thus, when isocurvature modes are inaccessible, we expect that decoherence will occur generically.³ We check this assertion explicitly in the next section for a simple two field model with two massive free fields. We find that this expectation is indeed correct for a relatively wide range of parameter values of the model.

IV. DECOHERENCE IN THE MODEL $V(\varphi, \chi) = \frac{1}{2}m^2\varphi^2 + \frac{1}{2}\mu^2\chi^2$

In this section we illustrate the role of isocurvature perturbations for decoherence in an inflationary model with $V(\varphi, \chi) = \frac{1}{2}m^2\varphi^2 + \frac{1}{2}\mu^2\chi^2$, a model which has been studied extensively in [23] (see also [18]). We first present exact numerical results for two cases: a) $\mu = 1.05m$ and b) $\mu = 10m$. We then derive an analytic result for the first case. We note that the first case is practically the same as single-field inflation as the trajectory in field space is practically straight. In the second case, the heavier field drops out of slow roll and start oscillating approximately 40 e-folds before inflation ends. However, as cosmologically observable scales exit the horizon while both fields are still in slow roll, the spectral index does not deviate significantly from unity and thus the model is observationally viable.

The wavefunction obtained from (32) explicitly reads:

$$i\hbar\partial_\eta\mathbf{B}_{11} = \hbar^2 (\mathbf{B}_{11}^2 + \mathbf{B}_{12}^2) + 2i\hbar\partial_\eta\theta\mathbf{B}_{12} - (k^2 + (aH)^2\Omega_{11}), \quad (41)$$

$$i\hbar\partial_\eta\mathbf{B}_{22} = \hbar^2 (\mathbf{B}_{22}^2 + \mathbf{B}_{12}^2) - 2i\hbar\partial_\eta\theta\mathbf{B}_{12} - (k^2 + (aH)^2\Omega_{22}), \quad (42)$$

$$i\hbar\partial_\eta\mathbf{B}_{12} = \hbar^2\mathbf{B}_{12}(\mathbf{B}_{11} + \mathbf{B}_{22}) + i\hbar(\mathbf{B}_{22} - \mathbf{B}_{11})\partial_\eta\theta - (aH)^2\Omega_{12}, \quad (43)$$

where

$$\Omega_{11} = -2 + \frac{1}{2H^2}(\mu^2 + m^2 - (\mu^2 - m^2)\cos 2\theta) - 5\epsilon - 2\epsilon(\epsilon + 2\eta^\parallel), \quad (44)$$

$$\Omega_{22} = -2 + \frac{1}{2H^2}(\mu^2 - m^2 + (\mu^2 + m^2)\cos 2\theta) + \epsilon, \quad (45)$$

$$\Omega_{12} = \Omega_{21} = \frac{\mu^2 - m^2}{2H^2}\sin 2\theta - 2\epsilon\eta^\perp \quad (46)$$

These equations are solved numerically with initial condition $\mathbf{B}_{11} = \mathbf{B}_{22} \simeq k/\hbar$, $\mathbf{B}_{12} = 0$. Figure 1 shows the evolution of

$$s_{\mathbf{k}} \equiv \frac{1}{2} \ln \frac{4C_{22}}{C_{11}} = \frac{1}{2} \ln \left[\frac{\Re[B_{11}]\Re[B_{11}] + \Im[B_{12}]^2}{\Re[B_{11}]\Re[B_{11}] - \Re[B_{12}]^2} \right], \quad (47)$$

which controls the amount of decoherence, for the two mass ratios $\mu/m = 1.05$ (blue, solid line) and $\mu/m = 10$ (black, solid line). This ratio is also the entropy per mode, see section V. Horizon exit occurs at 5 and 3 e-folds respectively

³ For the behavior of real and imaginary parts relating to inflationary perturbations see [4] and the next section




FIG. 1: The entanglement entropy per mode $s_{\mathbf{k}}$ plotted for the cases $\mu = 1.05m$ (numeric: solid, blue line; analytic: dashed-dotted, red line) and $\mu = 10m$ (numeric: solid, black line). For the first case, the analytic and numerical results are hardly distinguishable. Horizon exit happens at 5 and 3 e-folds respectively for the two cases and a few e-folds later the density matrix has decohered. For similar masses the entropy grows linearly with the number of e-folds until the end of inflation. For dissimilar masses, the entropy stops growing when the heavy field starts oscillating and remains constant on average, becoming zero for negligible time intervals.

for the chosen mode and inflation finishes at 60 e-folds. We see that $s_{\mathbf{k}}$ becomes larger than unity a few e-folds after horizon exit and continues to grow linearly with the number of e-fold causing the density matrix to decohere. For $\mu/m = 1.05$ this growth continues until the end of inflation. For the larger mass ratio it stops once the heavier field drops out of slow roll and starts oscillating, but s retains its average value until the end of inflation (it becomes zero for very short intervals and is quickly restored). Thus, in both cases decoherence is established a few e-folds after horizon crossing and is retained to the end of inflation.

The period of oscillations in figure 1 represents the Poincaré recurrence time for our system. The recurrence time is short because for each mode we have only two degrees of freedom. In models with several scalar fields however, we expect that the Poincaré recurrence time grows exponentially. Hence the brief recoveries of coherence we see in figure 1 should be a specific feature of the two scalar field model under consideration.

The case of almost equal masses can also be treated analytically where both fields exhibit slow-roll:

$$\frac{d\phi}{dN} \simeq \frac{\partial_{\phi} V}{3H^2}, \quad \frac{d\chi}{dN} \simeq \frac{\partial_{\chi} V}{3H^2}, \quad H^2 \simeq \frac{1}{3M_p^2} V \quad (48)$$

with $dN = -Hdt = -aHd\eta$, $M_p^2 = (8\pi G)^{-1}$, $H = \dot{a}/a$ the (almost constant) inflationary expansion rate of the universe. Denoting the number of e-folds N as

$$a(N) = a_{\star} e^{N_{\star} - N}, \quad (49)$$

we have that

$$\varphi = 2M_p \sqrt{N} \cos \omega \quad (50)$$

$$\chi = 2M_p \sqrt{N} \sin \omega. \quad (51)$$

As N decreases inflation proceeds, ending when $N \simeq 1$. The angular variable ω evolves according to

$$\frac{d\omega}{dN} = \frac{\mu^2 - m^2}{6H^2} \sin 2\omega. \quad (52)$$

Slow roll implies that

$$\tan \theta = \frac{\mu^2}{m^2} \tan \omega, \quad (53)$$

which gives

$$\frac{d\theta}{dN} = \frac{\mu^2 - m^2}{6H^2} \sin 2\theta. \quad (54)$$

The above expression can be easily integrated (see [23] for details). The expansion rate H is then given by

$$H^2 = \frac{1}{3}N (\mu^2 + m^2 - (\mu^2 - m^2) \cos 2\omega). \quad (55)$$

Furthermore, we are going to need an expression for the slow roll parameter

$$\epsilon = -\dot{H}/H^2 = \frac{1}{2M_p^2} \left[(d\varphi/dN)^2 + (d\chi/dN)^2 \right]. \quad (56)$$

We have

$$\epsilon = \frac{1}{N} \frac{m^4 + \mu^4 - (\mu^4 - m^4) \cos 2\omega}{[\mu^2 + m^2 - (\mu^2 - m^2) \cos 2\omega]^2}. \quad (57)$$

Let us first focus on subhorizon scales, where the $\partial_\eta \theta$ terms are not important; we have

$$i\hbar \partial_\eta \mathbf{B}_{11} = \hbar^2 (\mathbf{B}_{11}^2 + \mathbf{B}_{12}^2) - k^2, \quad (58)$$

$$i\hbar \partial_\eta \mathbf{B}_{12} = \hbar^2 2\mathbf{B}_{12}\mathbf{B}_{11} - (aH)^2 \Omega_{12}, \quad (59)$$

with solution $\mathbf{B}_{11} = \mathbf{B}_{22} \simeq k/\hbar$, $B_{12} = 0$, which is the standard Minkowski vacuum for two decoupled degrees of freedom.

On superhorizon scales, an analytic solution to (41) - (43) can be obtained perturbatively if $\partial_\eta \theta$ is treated as a small parameter. This will be true if

$$\frac{\mu^2 - m^2}{m^2} \ll 1. \quad (60)$$

To 0^{th} order in $\partial_\eta \theta$ we have

$$i\hbar \partial_\eta \mathbf{B}_{11} \simeq \hbar^2 (\mathbf{B}_{11}^2 + \mathbf{B}_{12}^2) - (k^2 + (aH)^2 \Omega_{11}), \quad (61)$$

$$i\hbar \partial_\eta \mathbf{B}_{22} \simeq \hbar^2 (\mathbf{B}_{22}^2 + \mathbf{B}_{12}^2) - (k^2 + (aH)^2 \Omega_{22}), \quad (62)$$

$$i\hbar \partial_\eta \mathbf{B}_{12} \simeq \hbar^2 \mathbf{B}_{12} (\mathbf{B}_{11} + \mathbf{B}_{22}). \quad (63)$$

Thus, $\mathbf{B}_{12} = 0$ and B_{11} , B_{12} can be obtained from the single field results (see (14)) where $B(\mathbf{k}, \eta) \equiv A(\mathbf{k}, \eta)/a^2$ and in the long wavelength limit

$$\lim_{k \ll aH} q_k \simeq \frac{e^{i\frac{\pi}{2}(\nu - \frac{1}{2})}}{\sqrt{2k\pi}} \Gamma(\nu) \left(-\frac{k\eta}{2} \right)^{-\nu + \frac{1}{2}}. \quad (64)$$

Using these, we find to leading order for the correlator

$$\mathbf{B}_{11} = \frac{aH}{\hbar} \left(\frac{2\pi}{\Gamma(\nu_1)^2} \left(\frac{k}{2aH} \right)^{2\nu_1} + i \left(\frac{3}{2} - \nu_1 \right) \right), \quad (65)$$

$$\mathbf{B}_{22} = \frac{aH}{\hbar} \left(\frac{2\pi}{\Gamma(\nu_2)^2} \left(\frac{k}{2aH} \right)^{2\nu_2} + i \left(\frac{3}{2} - \nu_2 \right) \right), \quad (66)$$

where

$$\nu_1 \simeq \frac{3}{2} + \frac{5}{3}\epsilon - \frac{1}{6H^2} (\mu^2 + m^2 - (\mu^2 - m^2) \cos 2\theta_0) \quad (67)$$

$$\nu_2 \simeq \frac{3}{2} - \frac{1}{3}\epsilon - \frac{1}{6H^2} (\mu^2 - m^2 + (\mu^2 + m^2) \cos 2\theta_0), \quad (68)$$




FIG. 2: *Left*: An illustration of the Wigner function before decoherence for a highly squeezed state describing long wavelength cosmological quantum fluctuations. The field amplitude (horizontal axis) is highly correlated with the field momentum (vertical axis). The area of the ellipse corresponds to minimum uncertainty and the entropy is zero. *Right*: The Wigner function after decoherence. The ellipse has increased its area leading to a non-vanishing value for the entropy. Note that the correlation between field amplitude and momentum has become less sharp but since $\Delta_{\pi(deco)}/\pi_{class} \ll 1$ even after decoherence, the prediction for acoustic oscillations in the CMB is not spoiled.

with $\tan \theta_0 = \frac{\mu^2}{m^2} \tan \omega_0$. Then, to first order in $\partial_\eta \theta$ we have

$$i\hbar \partial_\eta B_{12} \simeq \hbar^2 B_{12} (B_{11} + B_{22}) + [i\hbar (B_{22} - B_{11}) + 3aH] \partial_\eta \theta. \quad (69)$$

Soon after horizon crossing, B_{11} and B_{22} become imaginary and B_{12} is given by

$$B_{12} = -\frac{i}{\hbar} \int_{N_*}^N e^{\int_{N_*}^N \alpha dN'} (3 - \beta) aH \frac{\mu^2 - m^2}{6H^2} \sin 2\theta d\bar{N}, \quad (70)$$

where we have taken N_* to be a time when the real parts of B_{11} and B_{22} have become negligible, and

$$\alpha = \frac{4}{3}\epsilon - \frac{\mu^2 + m^2 \cos 2\theta_0}{3H^2} = \frac{\gamma}{N}, \quad (71)$$

$$\beta = 2\epsilon + \frac{\mu^2 \cos 2\theta_0 - m^2}{3H^2} \cos 2\theta_0 = \frac{\delta}{N}. \quad (72)$$

Using (55) and (57), and neglecting the dependence of θ on N in (70) we find

$$B_{12}(N) \simeq \frac{i}{\hbar} \frac{\mu^2 - m^2}{2\sqrt{3}} \frac{\sin 2\theta_0}{\sqrt{\mu^2 + m^2 - (\mu^2 - m^2) \cos 2\omega_0}} a(N) I(N), \quad (73)$$

where

$$\begin{aligned} I(N) &\equiv N^\gamma \int_N^{N_*} \left(3 - \frac{\delta}{s}\right) \frac{e^{N-s}}{s^{\gamma+\frac{1}{2}}} ds \\ &= N^\gamma e^N \left(\delta \Gamma\left(-\frac{1}{2} - \gamma, N_*\right) - 3 \Gamma\left(\frac{1}{2} - \gamma, N_*\right) - \delta \Gamma\left(-\frac{1}{2} - \gamma, N\right) + 3 \Gamma\left(\frac{1}{2} - \gamma, N\right) \right). \end{aligned} \quad (74)$$

As seen in figure 1, the above formulae reproduce the relevant numerical results with high accuracy.

V. ENTANGLEMENT ENTROPY

We now discuss how decoherence can be quantified by the amount of entanglement entropy for cosmological perturbations which is generated after tracing out one of the degrees of freedom. The analysis in [24, 25] implies that the

entropy of a gaussian quantum state can be expressed in terms of the product of the field amplitude and momentum variances (Δ_q^2 , Δ_π^2) as

$$S = -\text{tr} \ln [2\Delta_q \Delta_\pi / \hbar], \quad (75)$$

with $S = 0$ for any minimum uncertainty state such as the state of cosmological perturbations. The variance of the momentum Δ_π is calculated from the Wigner function $W(q, \pi)$ corresponding to the density matrix:

$$W(q, \pi) = \int d\pi d\pi^* \rho e^{-i\Delta\pi^* - i\Delta^* \pi}, \quad (76)$$

which for a gaussian state reads

$$W(q, \pi) \propto \exp \left(-\mathbf{q} \frac{1}{\Delta_q^2} \mathbf{q}^* - (\pi - \pi_{cl}) \frac{1}{\Delta_\pi^2} (\pi - \pi_{cl})^* \right). \quad (77)$$

The classical momentum associated with a value for $\mathbf{q}_1$ is

$$\pi_{1cl} = -\mathbf{q}_1 \mathbf{C}_{12} \hat{h}. \quad (78)$$

Note that the existence of a classical momentum, corresponding to a Wigner ellipse rotated in phase space (figure 2), does not alter the variances of $\mathbf{q}$ and π and equation (75) always holds as can be checked explicitly.

Before decoherence, $\Delta_{q_1}^2 = 1/\Re[\mathbf{B}_{11}]$ and $\Delta_{\pi_1}^2 = \hbar^2 \Re[\mathbf{B}_{11}]/4$ such that the state (15) corresponds to the minimum uncertainty state with vanishing entropy. Calculating the entanglement entropy corresponding to the reduced density matrix (37), we get for the decohered adiabatic perturbation

$$S = V \int \frac{d^3 k}{(2\pi)^3} s_{\mathbf{k}} = V \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2} \ln \left[\frac{4\mathbf{C}_{22}}{\mathbf{C}_{11}} \right]. \quad (79)$$

Thus, the entropy is characterized by the same quantity (47) which controls decoherence and figure 1 also shows the evolution of the entropy for the cases of the two mass ratios.

Equation (79) establishes a link between the entropy of cosmological perturbations and decoherence; the entropy (79) corresponds to the entanglement entropy which equals (minus) the information stored in correlations between the adiabatic and isocurvature perturbations. Therefore, it conforms with the standard definition of the entropy of quantum systems: tracing out the isocurvature perturbation (the unobservable “environment”) generates the entropy of the adiabatic perturbation (the system).

The effect of decoherence on the Wigner function is illustrated in figure 1. As mentioned above, prior to decoherence the state is described by the squeezed ellipse, the area of which is dictated by the minimum uncertainty. After decoherence, the variance Δ_{q_1} remains to a good accuracy unchanged but the variance in the momentum has increased to $\Delta_{\pi_1(dec)}^2 = 2\mathbf{C}_{22}$. This is opposite to the standard treatment of inflationary perturbations where the extremely squeezed nature of the state is taken to imply an exact classical correlation between the mode amplitude, drawn from a gaussian distribution, and the momentum. Decoherence implies that the momentum should also be drawn from a distribution with a dispersion larger than that dictated by the minimum quantum uncertainty.

The existence of acoustic peaks in the CMB requires that the field amplitude and the corresponding momentum exhibit a high degree of classical correlation as was already pointed out in [26]. Assuming $\mathbf{q}_1 \sim \Delta \mathbf{q}_1$, we find that the existence of such a classical correlation would require the variance of the momentum to be much smaller than the corresponding classical value, i.e.

$$\frac{\Delta_{\pi_1(dec)}}{|\pi_{1cl}|} = \frac{\sqrt{\mathbf{C}_{11}\mathbf{C}_{22}}}{|\mathbf{C}_{12}|} \ll 1. \quad (80)$$

This condition is satisfied for both cases as can be seen in figure 3. The second case exhibits an almost perfect degree of classical correlation by the time inflation ends and the oscillations of the heavier field have decayed. This behavior parallels that of the single field case, where the deviation from a perfect classical correlation of the field amplitude and the corresponding momentum is exponentially small [4]. The first case also exhibits correlation between field amplitude and momentum but which is significantly lower; the variance in the momentum is about 10^{-4} of the classical momentum value. In principle, such an effect will tend to wash out the CMB acoustic peaks slightly. Although it seems unlikely that such a small relative variance in the momentum would have an observable effect on the CMB, it may be interesting to examine its consequences.




FIG. 3: The classical correlation between field amplitude and momentum as quantified by the ratio $\ln \frac{\sqrt{C_{11}C_{22}}}{C_{12}}$ for $\mu = 1.05m$ (left) and $\mu = 10m$ (right).

VI. DISCUSSION

Let us close with a brief summary of the arguments presented in this paper. We discussed the possible relevance of isocurvature perturbations for the decoherence of cosmological curvature perturbations generated during inflation. We have argued that $q_1 - \Phi - q_2$ can play for cosmological perturbations the role of the triptych “system” - “measuring apparatus” - “environment” which is central to considerations of decoherence. The crucial ingredient is the appearance of a non-zero imaginary component B_{12} on superhorizon scales which results in $C_{11}/C_{22} \ll 1$ exponentially. We expect this phenomenon to be generic in all multi-field models since, in general, the imaginary parts are exponentially dominant over the real parts. We explicitly checked a free two-field model where no explicit couplings between the two fields were postulated. Nevertheless, adiabatic and isocurvature perturbations are generically coupled through gravitational interactions, implying that decoherence of this sort will be relevant in general. Since this decoherence is very effective, we do not expect the inclusion of non-linear interactions to affect this result. Furthermore, even if the entropy - and the momentum variance - increases, the strong classical correlation between field amplitude and momentum, responsible for the appearance of the CMB acoustic peaks, persists.

VII. ACKNOWLEDGMENTS

This research was supported by EU-RTN network MRTN-CT-2004-005104.

-
- [1] V. F. Mukhanov & G. V. Chibisov, JETP Lett. **33**, 10 (1981)
 - S. W. Hawking, Phys. Lett. B **115**, 295 (1982)
 - A. A. Starobinsky, Phys. Lett. B **117**, 175 (1982)
 - A. Guth & S. -Y Pi, Phys. Rev. Lett. **49**, 1110 (1982)
 - J. M. Bardeen, P. J. Steinhardt & M. S. Turner, Phys. Rev. D **28**, 679 (1983)
 - [2] D. N. Spergel *et al.*, [astro-ph/0603449].
 - [3] A. H. Guth and S. Y. Pi, Phys. Rev. D **32** (1985) 1899.
 - [4] D. Polarski and A. Starobinsky Class. Quant. Grav. **13** (1996) 377 [gr-qc/9504030]
 - [5] F. C. Lombardo, F. D. Mazzitelli and D. Monteoliva, Phys. Rev. D **62** (2000) 045016 [hep-ph/9912448]
 - [6] W. H. Zurek, Rev. Mod. Phys. **75** (2003) 715; For a short introduction see also Phys. Today, **44** (1991) 36
 - [7] E. Joos *et al.*, “Decoherence and the appearance of a classical world in quantum theory”, Springer-Verlag (2003)
 - [8] R. H. Brandenberger, R. Laflamme and M. Mijic, Mod. Phys. Lett. **A5** (1990) 2311
 - [9] E. Calzetta and B. L. Hu, Phys. Rev. D **52** (1995) 6770 [gr-qc/9505046]
 - [10] F. Lombardo and F. Mazzitelli, Phys. Rev. D **53** (1996) 2001 [hep-th/9508052]
 - [11] A. Matacz, Phys. Rev. D **55** (1997) 1860 [gr-qc/9604022]
 - [12] D. Campo and R. Parentani, Phys. Rev. D **74** (2006) 025001 [astro-ph/0505376]
 - [13] P. Martineau, [astro-ph/0601134]
 - [14] C. P. Burgess, R. Holman and D. Hoover, [astro-ph/0601646].
 - [15] C. Kiefer, I. Lohmar, D. Polarski and A. A. Starobinsky, [astro-ph/0610700].
 - [16] F. C. Lombardo, D. L. Nasir, Phys. Rev. D **72** (2005) 063506, [gr-qc/0506051]
 - [17] A. Albrecht, P. Ferreira, M. Joyce and T. Prokopec, Phys. Rev. D **50** (1994) 4807 [astro-ph/9303001].

- [18] S. G. Nibbelink and B. J. W. van Tent, *Class. Quant. Grav.* **19** (2002) 613 [hep-ph/0107272]
- [19] C. Gordon, D. Wands, B. A. Bassett, and R. Maartens, *Phys. Rev.* **D63** (2001) 023506 [astro-ph/0009131]
- [20] A. R. Liddle and D. H. Lyth, *Cosmological inflation and large-scale structure*, Cambridge Univ. Pr., UK: (2000) 400 p
- [21] D. H. Lyth and D. Wands, *Phys. Lett. B* **524** (2002) 5 [hep-ph/0110002].
- [22] K. Enqvist and M. S. Sloth, *Nucl. Phys. B* **626** (2002) 395 [hep-ph/0109214].
- [23] D. Polarski and A. A. Starobinsky, *Nucl. Phys. B* **385**, (1992) 623 ; *Phys. Rev. D* **50**, (1994) 6123 [astro-ph/9404061]
- [24] R. H. Brandenberger, T. Prokopec and V. F. Mukhanov, *Phys. Rev. D* **48** (1993) 2443 [gr-qc/9208009].
- [25] T. Prokopec, *Class. Quant. Grav.* **10** (1993) 2295.
- [26] C. Kiefer, D. Polarski, A. Starobinsky, *Phys. Rev. D.* **62**(2000) [gr-qc/9910065]